

Application of the collective model to determine some rotational bands of ^{239}U nucleus

Nguyen An Son¹, Le Viet Huy¹, Pham Ngoc Son²

Abstract – ^{238}U material is component in fuels of nuclear reactor core. Understanding properties and structure of ^{238}U nucleus is necessary before simulating and designing nuclear reactor. Besides that, the study of nuclear reaction is necessary to identify the specific characteristics of nucleus, it is the most effective experimental method up to now. However, in order to explain the properties of nuclear structure, in addition to study of the nuclear reaction, nuclear structure models and its theory must be used. There are many nuclear structure models to solve those properties of nucleus. This paper presents application of the Collective Model to determine some rotational bands of ^{239}U nucleus, using Prompt gamma neutron activation analysis method (PGNAA). Experiment is performed at channel No.2 of Dalat Research Reactor (DRR), using Filtered Thermal Neutron Beam and Compton Suppression Spectroscopy with High – Purity Germanium detector (HPGe). The results have found 11 rotational bands of ^{239}U nucleus. This work is very necessary for the research of nuclear structure which controls material technology by itself.

Index Terms – Collective model, ^{239}U , rotational bands

1 INTRODUCTION

Collective Model was developed in the 1950s by Reynolds, A. Bohr and Mottelson, Hill and Wheele [1]. The Collective Model emphasizes the coherent behavior of all nucleons in heavy nuclei. The spherical symmetric potential of the nucleus with full shell is exceptionally stable to the effects of additional nucleons. Therefore, it still remains spherical symmetric form. The excited state of a nucleus is defined by single-particle levels in the spherical symmetric potential and the quadrupole field of a spherical symmetric nucleus.

By the increase of external nucleons of the full shell of nucleus, the individual motion effect of nucleons on the potential field increases and the centrifugal pressure of nucleons appears.

Collective motion increases rapidly and impact the core of full shell nucleus leading to the decrease of the potential field stability, it means that the nucleus has spherical asymmetric form. According to quantum mechanics, spherical asymmetric nuclei can rotate.

The Collective Model has been very successful in describing variety of nuclear properties, especially energy levels in nuclei that the Shell Model and the Liquid Drop Model cannot be applied. These energy levels show the characteristics of rotating or vibrating systems. The properties of these nuclei, including excited state energies, angular momentum, magnetic moments, nuclear shapes, etc. can be understood by using the Collective Model.

In 1969, Larry Shelton Varnell [2] applied the Collective Model to determine the rotational band of some deformed nuclei, using the vacuum chamber of the Si(Li) electron spectrometer with Lithium Drifted Silicon detector. The result had determined 12 rotational bands of ^{152}Sm , 14 rotational bands of ^{154}Gd , and 12 rotational bands of ^{166}Er .


So far, there are many researches on ^{239}U nucleus. In 1959, the smoothed gamma-ray spectrum due to neutron capture by ^{238}U obtained by Campion et al [3]. The gamma rays in the energy region between 0.14 MeV and 3.4 MeV were examined with a crystal spectrometer. The gamma-rays in the 3.4 MeV to 4.2 MeV range were studied using a pair spectrometer with a resolution of about 1%. In the 1972s, Booth et al [4] had found 21 discrete energy levels and the spin-parity assignments of the excited ^{239}U nucleus up to 0.950 MeV. In the 1970s, John et al [5] studied about neutron capture gamma radiation from neutron capture in ^{238}U . The results had found 16 energy gamma-rays from 1 MeV to 4.75 MeV, with the intensity was reported on a number per one hundred capture events, etc. But there have

Received: 17-07-2017, Accepted: 14-08-2017, Published: 10-8-2018

Author: Nguyen An Son, Le Viet Huy- Da Lat University (e-mail: sonna@dlu.edu.vn) Pham Ngoc Son - Nuclear Research Institute

no research before about the rotational band of ^{239}U nucleus.

Prompt gamma neutron activation analysis (PGNAA) is a rapid, nondestructive sample technique which is very effective in nuclear structure research [6]. ^{238}U exists in nature, its abundance is about 99.27%, which is component in fuels of nuclear reactor core. ^{238}U is heavy even-even nucleus which has 92 protons and 146 neutrons. ^{239}U is heavy even-odd nucleus which has 92 protons, 146 neutrons and 1 added neutron. In this experiment, PGNAA method is used to acquire the prompt gamma-rays emitted from ^{238}U (n, γ) ^{239}U reaction.



Where n is incident neutron, ^{238}U is the target nucleus, $({}^{239}\text{U})^*$ is compound nucleus, $({}^{239}\text{U})$ is product nucleus and γ_{prompt} is prompt gamma-rays.

2 THEORY AND EQUIPMENTS

Theory

The Nilsson model is a shell model for a deformed nucleus. It provides a description of single-particle motion in a spherical asymmetric potential. An appropriate single-particle Hamiltonian for a nucleus with the symmetry axis z is given by [7]:

$$H = T + V = \frac{p^2}{2m} + \frac{1}{2}m[\omega_x(x^2 + y^2) + \omega_z z^2] + Cls + Dl^2 \quad (1)$$

where ω_x , ω_y and ω_z are one-dimensional oscillator frequencies in the x , y , and z direction. C and D are constant. The l^2 and ls terms ensure the proper order and energies of the single-particle levels in the spherical limit.

$$\omega_x^2 = \omega_y^2 = \omega_0^2 \left(1 + \frac{2}{3}\delta\right);$$

$$\omega_z^2 = \omega_0^2 \left(1 - \frac{4}{3}\delta\right);$$

$$\omega_0 = \left(1 - \frac{4}{3}\delta^2 - \frac{16}{27}\delta^3\right)^{\frac{1}{6}} = \text{constant}$$

where ω_0 is the oscillator frequency in the spherical potential. It is assumed that the nuclear volume remains constant as a function of ω_0 . The rotational Hamiltonian is of the form [7]:

$$H_{\text{rot}} = \frac{\Omega^2}{2J_{\text{eff}}} = \frac{I^2 - K^2}{2J_{\text{eff}}} = \frac{\hbar^2}{2J_{\text{eff}}}[I(I+1) - K(K+1)] \quad (2)$$

where I is a total angular momentum, Ω is rotational angular momentum and K is the spin of nucleus.

$$\vec{I} = \vec{K} + \vec{\Omega};$$

$$I^2 = \hbar^2 I(I+1); K^2 = \hbar^2 K(K+1)$$

J_{eff} is the effective moment of inertia,

$$J_{\text{eff}} \approx J_0 \left(\frac{\Delta R}{R}\right)^2$$

where J_0 is the moment of the inertia of nucleus, R is radius of nucleus and ΔR is a deformation parameter of nucleus.

In deformed even-even nuclei, the spin of nucleus is in the ground state ($K=0$), equation (2) can be written [1]:

$$E_{\text{rot}} = \frac{\hbar^2}{2J_{\text{eff}}} I(I+1)$$

where $I = 0, 2, 4, 6, \dots$ for positive parity states ($\pi = +1$) and $I = 1, 3, 5, 7, \dots$ for negative parity states ($\pi = -1$).

In deformed odd-A and odd-odd nuclei ($K \neq 0$), equation (2) can be written [1]:

$$E_{\text{rot}} = \frac{\hbar^2}{2J_{\text{eff}}}[I(I+1) - K(K+1)]$$

where $I = K, K+1, K+2, K+3, \dots$

^{239}U nucleus has spin $K = 5/2$ and $I = K, K+1, K+2, \dots = 5/2, 7/2, 9/2, \dots$. Thus, rotational energies are:

$$E_0 = 0; E_1 = \frac{7\hbar^2}{2J_{\text{eff}}}; E_2 = \frac{16\hbar^2}{2J_{\text{eff}}}; E_3 = \frac{27\hbar^2}{2J_{\text{eff}}};$$

$$E_4 = \frac{40\hbar^2}{2J_{\text{eff}}}; E_5 = \frac{55\hbar^2}{2J_{\text{eff}}}; E_6 = \frac{72\hbar^2}{2J_{\text{eff}}}; \dots$$

Then, the ratio between rotational energies is given by:

$$E_1/E_2/E_3/E_4/E_5/E_6/\dots = 1/16/27/40/55/72/\dots$$

Equipments

The experiment is performed at channel No.2 of DRR, which using Filtered Thermal Neutron Beam, and HPGe detector with PGNAA method. Configuration of the system is shown in Fig. 1.




Fig. 1. Configuration of the acquisition system at channel No.2 of DRR

The thermal neutron flux at the sample position is 1.6×10^6 n/cm²xs, and the Cd ratio is 420 [8]. Inside the channel No.2, a chamber with the internal high density polyethylene (HDPE) is set up, it also has 5% Li to shield the scattered neutrons. In the mid-core of this chamber, a holder is made of PTFE (Teflon plastic) material which fixed the sample during the acquisition process.

Due to the large number of gamma-rays incident on the main detector, the Compton continuum is a significant hindrance for low background. The Compton continuum causes the difficult search of low-intensity peaks and increases the uncertainty of the measured activities. Therefore, a Compton suppression spectroscopy has been set-up and installed at DRR 500 kW for neutron activation analysis and nuclear data measurement. The central detector is a GR7023 Canberra n-type coaxial HPGe detector. Its FWHM is 2.36 keV for the 1.33 MeV of ⁶⁰Co peak. The relative efficiency is 72%. There are 12 Bismuth Germanium (BGO) guard detectors shielded by a lead of 10 cm thickness. A lead-stepped collimator is located in the front of the opening of the guard detectors. The length and inner diameter of the lead collimator are 180 mm and 40 mm, respectively. The reduction of the Compton continuum has been achieved by surrounding the HPGe detector with the BGO detectors whose signals are used for the anti-coincidence gating in the analog-to-digital converter (ADC). The Compton continuum is reduced about 1.5 to 2 times, up to 1 MeV region of energy [9]. The detectors and shielding system are configured as Fig. 2.




Fig. 2. The back and the cross-sectional view of the detectors and shielding system

The electronic modules are manufactured by Canberra except the high voltage module for BGO detectors, which were produced by Fast Comtec. They include 2026 main amplifiers (AMP), 3106D high voltage power supply, multiport II with ADC 16K and multichannel analyzer (MCA), using the Genie 2000 software. Its configuration is shown in Fig. 3.





Fig. 3. The block schema of the gamma acquisition system

²³⁸U natural metal is used. Its diameter, thickness and weight are 1.2 cm, 0.5 cm and 23.68586 g respectively. Geometric form of ²³⁸U sample is cylinder form, which is shown in Fig. 4.

The ²³⁸U sample is placed in the holder at the irradiation position, the angle between the neutron flux and the sample is 45°, the distance from the sample to the detector is 38.5 cm.



Fig. 4. Geometric form of 238U



Energy Calibration: $y = 0.5101x - 0.354$
where x is the channel number and y is the gamma energy (keV)

Fig. 5. Prompt gamma spectrum of ^{239}U

3 RESULTS AND DISCUSSION

The acquisition time of background spectrum is 62,465 seconds and ^{239}U spectrum is 86,492 seconds.

Prompt gamma spectrum of ^{239}U acquired at channel No.2 of DRR after eliminating the effect

of background is shown in Fig. 5. The statistical count of the spectrum is 1.92×10^8 counts.

Experimental data are shown in Table 1. There are 36 prompt gamma-rays emitted from ^{238}U (n, γ) ^{239}U reaction. Determination of rotational bands of ^{239}U is calculated by Equation (5) and (6). Results compared between experimental data and theoretical calculation are shown in Table 2.

Table 1. Energy and Intensity of prompt gamma-rays emitted from ^{238}U (n, γ) ^{239}U reaction

No.	Energy (keV)	Intensity (%)	No.	Energy (keV)	Intensity (%)	No.	Energy (keV)	Intensity (%)
1	96.74	0.7403	13	431.34	1.9458	25	833.25	4.0339
2	111.20	0.4581	14	456.27	2.0985	26	847.22	5.6266
3	139.79	0.0057	15	481.20	2.2686	27	867.66	1.8530
4	160.07	0.2853	16	510.62	9.2938	28	915.03	3.7763
5	174.53	0.2840	17	538.55	2.9969	29	974.87	3.9210
6	197.96	0.0382	18	553.01	3.5011	30	1097.54	3.4827
7	252.32	0.6360	19	595.89	1.6280	31	1137.44	3.4598
8	277.75	0.9082	20	609.36	3.8907	32	1202.76	3.3756
9	296.20	1.0473	21	695.63	3.7557	33	1221.71	5.1242
10	326.12	0.9353	22	707.09	3.3245	34	1279.05	4.6383
11	352.05	1.4591	23	777.90	6.4912	35	1295.51	2.9494
12	398.43	1.7781	24	810.82	3.3157	36	1311.97	4.6733

Table 2. Results compared between experimental data and theoretical calculation

No.	E_γ experimental (keV)	(E_i/E_1) experimental (keV)	(E_i/E_1) theory (keV)	No.	E_γ experimental (keV)	(E_i/E_1) experimental (keV)	(E_i/E_1) theory (keV)
1	(42.53)	1.00	1.00	7	553.01	13.00	13.00
2	96.74	2.27	2.29	8	695.63	16.36	16.00
3	160.07	3.76	3.86	9	810.82	19.06	19.26
4	252.32	5.93	5.71	10	974.87	22.92	22.86
5	326.12	7.67	7.86	11	1137.44	26.74	26.71
6	431.34	10.14	10.26	12	1311.97	30.85	30.86

*Note: (42,53) keV is taken from Nuclear Data Center [10]

Results in Table 2 show that ^{239}U nucleus has 11 rotational bands, which are 96.74 keV; 160.07 keV; 252.32 keV; 326.12 keV; 431.34 keV; 533.01 keV; 695.63 keV; 810.82 keV; 974.87 keV; 1137.44 keV and 1311.97 keV. Among 36 energy peaks from the prompt gamma spectrum of ^{239}U nucleus, 25 another peaks are from ^{235}U (n, γ) ^{236}U reaction (abundance of ^{235}U nucleus in the sample is about 0.73%) and from the background spectrum. However, we can't find the 42.53 keV peak from the spectrum which is the first excited state of ^{239}U nucleus [10]. It's the limitation of experimental procedure.

However the determination of 11 peak energies of the experimental spectrum is very closed to the theoretical calculation of rotational bands by using Equation (6). Therefore we can conclude that the 42.53 keV energy is a part of rotational bands.

4. CONCLUSION

From prompt gamma spectra acquired at the channel No.2 of DRR using application of Collective Model in nuclear structure research, some rotational bands of ^{239}U deformed nucleus are identified.

The results are quite relevant to the theory of the Collective Model when studying about the heavy nucleus, which has large different between the neutron and proton numbers. These results have

shown that the ^{239}U deformed nucleus has spherical asymmetric structure.

REFERENCES

- [1]. A. Bohr, B.R. Mottelson, Nuclear Structure, *World Scientific Publishing*, pp. 24, 33, 335, 1998.
- [2]. L.S. Varnell, Beta, Gamma Vibrational bands in Deformed Nuclei, *California Institute of Technology*, May 19, 1969.
- [3]. P.J. Campion, J.W. Knowles, G. Manning, G. A. Bartholomew, *Canadian Journal of Physics*, 1959.
- [4]. R.S. Booth, J.E. White, S.K. Penny, K.J. Yost, *Nuclear Science and Engineering*, 1972.
- [5]. J. Joh, V.J. Orphan, Gamma Rays From Resonant Capture of Neutrons in ^{238}U , GA-10186, Gulf General Atomic (1970).
- [6]. Z.B. Alfassi, Prompt Gamma Neutron Activation Analysis with Reactor Neutrons, pp. 59, 1995.
- [7]. R.F. Casten, Nuclear Structure from a Simple Perspective, Oxford University Press, pp 167, 256, 257, 1990.
- [8]. P.N. Son, "Phát triển dòng neutron phân loại trên kênh ngang số 2 của Lò phản ứng hạt nhân Đà Lạt", *Báo cáo Tổng kết đề tài nghiên cứu khoa học cấp Bộ*, 2011.
- [9]. N.X. Hai, N.N. Dien, P.D. Khang, V.H. Tan, N.D. Hoa, "A simple configuration setup for Compton Suppression Spectroscopy", Cornell University Library 2013.
- [10]. National Nuclear Data Center: <https://www.nndc.bnl.gov/chart/reColor.jsp?newColor=es>

Ứng dụng mẫu hạt nhân suy rộng trong việc xác định một số phổ quay của hạt nhân ^{239}U

Nguyễn An Sơn¹, Lê Viết Huy¹, Phạm Ngọc Sơn²

¹Trường Đại học Đà Lạt

²Viện Nghiên cứu Hạt nhân, Lâm Đồng, Việt nam

Tác giả liên hệ: sonna@dlu.edu.vn

Ngày nhận bản thảo: 17-07-2017, ngày chấp nhận đăng: 14-08-2017, ngày đăng: 10-08-2018

Tóm tắt – Vật liệu ^{238}U là thành phần trong nhiên liệu của lõi lò phản ứng hạt nhân. Việc tìm hiểu tính chất, cấu trúc của hạt nhân ^{238}U là cần thiết trước khi muốn mô phỏng, thiết kế lò phản ứng. Bên cạnh đó, nghiên cứu phản ứng hạt nhân là cần thiết trong việc xác định tính chất đặc thù của từng hạt nhân và là phương pháp thực nghiệm hữu hiệu cho đến ngày nay. Tuy nhiên, để giải thích các tính chất về cấu trúc hạt nhân, ngoài việc nghiên cứu phản ứng thì phải sử dụng các mẫu cấu trúc hạt nhân để làm sáng tỏ vấn đề. Có nhiều mẫu cấu trúc hạt nhân khác nhau để giải quyết cho bài toán này.

Bài báo trình bày ứng dụng mẫu suy rộng trong việc xác định một số phổ quay của hạt nhân ^{239}U , sử dụng phương pháp phân tích kích hoạt neutron đo gamma tức thời (PGNAA). Thục nghiệm được tiến hành tại kênh ngang số 2 của Lò phản ứng hạt nhân Đà Lạt (DRR), sử dụng dòng neutron phin lọc đơn năng và hệ phổ kế triệt Compton với đầu dò bán dẫn HPGe. Kết quả đã xác định được 11 phổ quay của hạt nhân ^{239}U . Đây là công việc thiết thực trong nghiên cứu cấu trúc hạt nhân và làm chủ công nghệ vật liệu.

Từ khóa – Mẫu suy rộng, ^{239}U , phổ quay